

Regression models and panel data

Part III: Panel data models

Rodolfo Metulini

✉ rmetulini@unisa.it

Department of Economics and Statistics (DISES) - University of Salerno

Outline

- 1 The one-way (error) component model
- 2 Ordinary least squares
- 3 Generalized least squares
- 4 Estimation comparison
- 5 Two ways error model
- 6 Advanced arguments
- 7 References

The one-way
(error)
component
model

Ordinary least
squares

Generalized
least squares

Estimation
comparison

Two ways
error model

Advanced
arguments

References

Further examples

- ① in an earnings equation in labor economics
$$y_{nt} = \alpha + x_{nt}^T \beta + \mu_n + v_{nt}$$
 y_{nt} measure earnings of the head of the household, x_{nt} contain a set of variables like experience, education, union membership, sex, race, etc. Note that μ_n is time-invariant and it accounts for any individual-specific effect that is not included in the regression. In this case we could think of it as the individual's unobserved ability. The remainder disturbance v_{nt} varies with individuals and time and can be thought of as the usual disturbance in the regression.
- ② for a production function utilizing data on firms across time, y_{nt} measure output and X_{nt} measure inputs. The unobservable firm-specific effects will be captured by the μ_n and we can think of these as the unobservable entrepreneurial or managerial skills of the firm's executives.

The model (i)

- For the observation of individual n at period t we can write the most simple version of the panel model as:

$$y_{nt} = \alpha + x_{nt}^T \beta + e_{nt} \quad (1)$$

or, equivalently:

$$y_{nt} = z_{nt}^T \gamma + e_{nt} \quad (2)$$

- The error component is the sum of two effects: μ_n , which is the effect for individual n ; v_{nt} is the residual (or idiosyncratic) effect: $e_{nt} = \mu_n + v_{nt}$

The model (ii)

- For the whole sample of $O = NT$ observations we have:

$$y = \alpha j + X\beta + e \quad (3)$$

or, equivalently:

$$y = Z\gamma + e \quad (4)$$

where y is a vector of length NT , X is a matrix of dimension $NT \times K$, Z is $NT \times (K + 1)$ and β is a vector of length K

The model (iii)

The one-way
(error)
component
model

Ordinary least
squares

Generalized
least squares

Estimation
comparison

Two ways
error model

Advanced
arguments

References

$$y = \begin{pmatrix} y_{11} \\ y_{12} \\ \dots \\ y_{1T} \\ y_{21} \\ y_{22} \\ \dots \\ y_{2T} \\ \dots \\ y_{N1} \\ y_{N2} \\ \dots \\ y_{NT} \end{pmatrix} \quad X = \begin{pmatrix} x_{11}^1 & x_{11}^2 & \dots & x_{11}^K \\ x_{12}^1 & x_{12}^2 & \dots & x_{12}^K \\ \dots & \dots & \dots & \dots \\ x_{1T}^1 & x_{1T}^2 & \dots & x_{1T}^K \\ x_{21}^1 & x_{21}^2 & \dots & x_{21}^K \\ x_{22}^1 & x_{22}^2 & \dots & x_{22}^K \\ \dots & \dots & \dots & \dots \\ x_{2T}^1 & x_{2T}^2 & \dots & x_{2T}^K \\ \dots & \dots & \dots & \dots \\ x_{N1}^1 & x_{N1}^2 & \dots & x_{N1}^K \\ x_{N2}^1 & x_{N2}^2 & \dots & x_{N2}^K \\ \dots & \dots & \dots & \dots \\ x_{NT}^1 & x_{NT}^2 & \dots & x_{NT}^K \end{pmatrix} \quad \beta = \begin{pmatrix} \beta_1 \\ \dots \\ \beta_K \end{pmatrix} \quad e = \begin{pmatrix} e_{11} \\ e_{12} \\ \dots \\ e_{1T} \\ e_{21} \\ e_{22} \\ \dots \\ e_{2T} \\ \dots \\ e_{N1} \\ e_{N2} \\ \dots \\ e_{NT} \end{pmatrix}$$

Transformation

- Panel data models
break the total variation in intra individual and inter individual
- The inter individual transformation (between) is obtained by post multiplying $B = I_N \otimes J_T / T$ with y and x
- For example, for one regressor, we have
 $(Bx)^T = (\bar{x}_{1\cdot}, \bar{x}_{1\cdot}, \dots, \bar{x}_{1\cdot}, \bar{x}_{2\cdot}, \bar{x}_{2\cdot}, \dots, \bar{x}_{2\cdot}, \dots, \bar{x}_{N\cdot}, \bar{x}_{N\cdot}, \dots, \bar{x}_{N\cdot})$
- The intra individual transformation (within) is obtained by post multiplying $W = I_{NT} - B$ with y and x
- For example, for one regressor, we have
 $(Wx)^T = (x_{11} - \bar{x}_{1\cdot}, x_{12} - \bar{x}_{1\cdot}, \dots, x_{1T} - \bar{x}_{1\cdot}, x_{21} - \bar{x}_{2\cdot}, x_{22} - \bar{x}_{2\cdot}, \dots, x_{2T} - \bar{x}_{2\cdot}, \dots, x_{N1} - \bar{x}_{N\cdot}, x_{N2} - \bar{x}_{N\cdot}, \dots, x_{NT} - \bar{x}_{N\cdot})$
- W and B are:
 - ① symmetric, so $B^T = B$ and $W^T = W$
 - ② idempotent, so $WxW = W$ and $BxB = B$
 - ③ They decompose a vector, so $Bxz + Wxz = z$
 - ④ orthogonal, so $W^TxB = 0$

Errors assumptions

- the errors, in matrix form, can be written as $e = (I_N \otimes J_T)\mu + v$
- The estimated model with estimated parameters $\hat{\gamma}^T = (\hat{\alpha}, \hat{\beta}^T)$ and the NT vector of residuals \hat{e} ($y = Z\hat{\gamma} + \hat{e}$) is:
 - $\hat{e} = e - Z(\hat{\gamma} - \gamma)$
- Assumptions:
 - ① the expected value of μ_n and v_{nt} is 0
 - ② the effects μ_n are mutually uncorrelated and homoschedastic
 - ③ the idiosyncratic terms v_{nt} are also mutually uncorrelated and homoschedastic
 - ④ v_{nt} and μ_n are uncorrelated each others
 - ⑤ the variance $E(e_{nt}^2) = \sigma_\mu^2 + \sigma_v^2$
 - ⑥ covariance $E(e_{nt}^2 e_{ns}^2), s \neq t = \sigma_\mu^2$
 - ⑦ covariance $E(e_{nt}^2 e_{mt}^2), n \neq m = 0$

The covariance matrix

- For a given individual n , the covar matrix for $e_n^T = (e_{n1}, e_{n2}, \dots, e_{nT})$ is

$$\Omega_{nn} = E(e_n e_n^T) = \sigma_v^2 I_T + \sigma_\mu^2 J_T \quad (5)$$

- for $n \neq m$ the covariance matrix is 0, given the assumptions of mutual uncorrelation. It follows that:
- The covariance matrix Ω is a N-block diagonal matrix, each one is in the form of equation 5:

$$\Omega = \sigma_v^2 I_{NT} + \sigma_\mu^2 (I_N \otimes J_T)$$

- Given B and W , $\Omega = \sigma_v^2 (B + W) + T \sigma_\mu^2 B = \sigma_v^2 W + \sigma_1^2 B$
- A desirable assumption is that both μ and v are uncorrelated with x , that is $E(\mu | x) = 0$ and $E(v | x) = 0$

OLS estimator(s)

- The variability in a panel has two sources:
 - ① between, or inter-individual, which is the variability of panel variables measured in individual means which is \bar{z}_n (or Bz)
 - ② within, or intra individual, which is the variability of panel variables measured in deviation from the means (i.e., $z_{nt} - \bar{z}_n$, or Wz)
- OLS method of estimation can be applied to 3 models: the *raw data* one, the *within* and the *between*

On raw data: pooling model

- The model to be estimated is $y = Z\gamma + e$
- OLS is based on minimizing the quantity $\hat{e}^T \hat{e}$
- the OLS estimator is

$$\hat{\gamma}_{OLS} = (Z^T Z)^{-1} Z^T y \quad (6)$$

while the difference between the vector of true parameters and estimated parameters is $\hat{\gamma}_{OLS} - \gamma = (Z^T Z)^{-1} Z^T e$

- About β , $\hat{\beta}_{OLS} = (X^T(I - \bar{J})X)^{-1} X^T(I - \bar{J})y$ with $E(\hat{\beta}) = \beta$ (unbiased) if the error e and X are uncorrelated
- $V(\hat{\gamma}_{OLS}) = (Z^T Z)^{-1} Z^T \Omega Z (Z^T Z)^{-1} \neq \sigma^2 (Z^T Z)^{-1}$
- So, OLS is unbiased and consistent if e and X are uncorrelated, but:
 - ① the expression for the variance is complex
 - ② OLS is not the BLUE: it exist at least one estimator which is unbiased and more efficient.

The between estimator

The one-way
(error)
component
model

Ordinary least
squares

Generalized
least squares

Estimation
comparison

Two ways
error model

Advanced
arguments

References

- Between estimator is an OLS applied to the between transformed model

$$By = BZ\gamma + Be = \alpha j + BX\beta + Be$$

- The variables (such as, e.g., gender, religion) in the model that do not exhibit intra-individual variation are **unaffected** by this transformation
- The NT observations become in fact N distinct observations with individual mean repeated T times each.
- $\hat{\beta}_B = (X^T \bar{B}X)^{-1} X^T \bar{B}y$, with $\bar{B} = B - \bar{J}$ the matrix that transform the variable in its mean into a deviation from the overall mean
- $V(\hat{\beta}_B) = \sigma_1^2 (X^T \bar{B}X)^{-1}$

The within estimator

- Within estimator is an OLS applied to the within transformed model (μ from $e = \mu + v$ wipes out)

$$Wy = W(\alpha j + X\beta + e) = WX\beta + Wv$$

- This transformation removes the vector of ones associated to the intercept α but also the matrix associated to the vectors of individual effects μ_n .
- It also **removes** regressors without intra-individual variation
- Applying OLS to the transformed model we have the within estimator

$$\hat{\beta}_W = (X^T W X)^{-1} X^T W y \quad (7)$$

with standard expression for the variance (no μ_n)

$$V(\hat{\beta}_W) = (X^T W X)^{-1} X^T W \Omega W X (X^T W X)^{-1} = \sigma_v^2 (X^T W X)^{-1}$$

- The within model is also called "fixed effects model" because individual effects are taken as **fixed parameters**

Least squares dummy variables (LSDV)

- The model can be written as:

$$y = X\beta + (I_N \otimes J_T)\mu + v$$

where μ is a N -length vector of parameters to be estimated (μ is no more stochastic)

- Therefore, there are $N + K$ parameters and the model is feasible only if N is not too large compare to T
- β coefficients estimated with within or LSDV are the same (Frisch-Waugh theorem)
- However, LSDV directly estimates the vector of parameters μ while the within just the β s
- Within estimates μ_n (if needed) in a separate step, by performing $\hat{\alpha}_n = \bar{y}_{n.} - \bar{x}_{n.}^T \hat{\beta}$, with $\hat{\mu}_n = \hat{\alpha}_n - \hat{\alpha}$ and $\hat{\alpha}$ is the intercept

Application

Application in R: Example 2.1 (Croissant, Millo)

GLS estimator (i)

- The within estimator is a regression on data that have been transformed so that individual effects vanish while LSDV estimates all the effects, but the risk is of having too many coefficients to be estimated
- On the contrary, GLS consider the individual effect as a random (stochastic) draw from a specific distribution (e.g., normal) and seeks to estimate the parameters (e.g. the variance) of this distribution
- In any case, the aim is to obtain an efficient estimate for the slopes (i.e., the β parameters)
- Assume $\mu_n \sim iid(0, \sigma_\mu^2)$, $v_{nt} \sim iid(0, \sigma_v^2)$ and the μ_n are independent of the v_{nt} .
- The random effects model is an appropriate specification if we randomly draw N individuals from a large populations where N is large. LSDV would lead to an enormous loss of degrees of freedom.

GLS estimator (ii)

- Assuming independence between the error terms μ_n and v_{nt} , the covariance matrix is

$$\Omega = E(e^T e) = Z_\mu E(\mu^T \mu) Z_\mu^T + E(v^T v) = \sigma_\mu^2 (I_N \otimes J_T) + \sigma_v^2 (I_N \otimes I_T)$$

- This implies a homoskedastic variance

$$\text{var}(e_{nt}) = \sigma_\mu^2 + \sigma_v^2, \forall n, t$$

- Ω is equicorrelated and block-diagonal and it exhibits serial correlation over time only between the same individual. In fact,

$$\text{cov}(e_{nt}, e_{ms}) = \begin{cases} \sigma_\mu^2 + \sigma_v^2 & \text{if } n = m, s = t \\ \sigma_\mu^2 & \text{if } n = m, s \neq t \\ 0 & \text{otherwise} \end{cases}$$

GLS estimator (iii)

$$\hat{\gamma}_{GLS} = (Z^T \Omega^{-1} Z)^{-1} (Z^T \Omega^{-1} y) \quad (8)$$

- $V(\hat{\gamma}_{GLS}) = (X^T \Omega^{-1} X)^{-1}$
- Ω is also linear combination of two well-known idempotent and orthogonal matrices: $\Omega^{-1} = \frac{1}{\sigma_1^2} B + \frac{1}{\sigma_v^2} W$
- Since the dimension of Ω depends on the sample, may be infeasible to estimate $\hat{\gamma}_{GLS}$ with eq. 8.
- An efficient way is to estimate OLS on pre transformed data.
- Let C such that $C^T C = \Omega^{-1}$, $\tilde{y} = Cy$, $\tilde{Z} = CZ$, then $\hat{\gamma} = (\tilde{Z}^T \tilde{Z})^{-1} (\tilde{Z}^T \tilde{y})$
- $C = \Omega^{-0.5} = \frac{1}{\sigma_1} B + \frac{1}{\sigma_v} W$, with σ_1 and σ_v that need to be estimated in advance

GLS estimator (iv)

- we can denote $\phi = \frac{\sigma_v}{\sigma_1}$.

It follows that $\theta = 1 - \phi = 1 - \sqrt{\frac{\sigma_v^2}{T\sigma_\mu^2 + \sigma_v^2}}$

- As Ω depends on W and B , it may be clear that GLS produces results in between the within (is equal to the within when σ_μ dominates, $\theta = 1$) and the pooled OLS (is equal to the pooled OLS when σ_v dominates, $\theta = 1$)

Application

Application in R: Example 2.2 (Croissant, Millo)

Comparison of the estimators

The one-way
(error)
component
model

Ordinary least
squares

Generalized
least squares

Estimation
comparison

Two ways
error model

Advanced
arguments

References

- We have four different estimators for the same model (the one-way error component model): the within and the between just use one source of variance (respectively, intra and inter-individual), the pooled OLS and the GLS use both sources.
- If the assumption on null correlation between the errors and the regressors is in place, all estimators are consistent and unbiased and they give similar results

Relations between estimators (i)

The one-way
(error)
component
model

Ordinary least
squares

Generalized
least squares

Estimation
comparison

Two ways
error model

Advanced
arguments

References

- OLS and GLS gives intermediate results between the *within* and the *between* estimators, as they use both sources of variance.
- It is in fact demonstrated that $\hat{\beta}_{GLS}$ can be expressed as a weighted average of the *within* and the *between*

$$\hat{\beta}_{GLS} = (X^T W X + \phi^2 X^T \bar{B} X)^{-1} (X^T W X \hat{\beta}_W + \phi^2 X^T \bar{B} X \hat{\beta}_B)$$

- A similar thing with $\hat{\beta}_{OLS}$, which is the GLS when $\phi = \phi^2 = 1$

$$\hat{\beta}_{OLS} = (X^T W X + X^T \bar{B} X)^{-1} (X^T W X \hat{\beta}_W + X^T \bar{B} X \hat{\beta}_B)$$

- For OLS the weights are exactly the shares of the intra- and inter-individual variance of the regressors.
- For GLS, the weights depends also on the variance of the errors.

Relations between estimators (ii)

The one-way
(error)
component
model

Ordinary least
squares

Generalized
least squares

Estimation
comparison

Two ways
error model

Advanced
arguments

References

- Because $\phi \leq 1$, GLS will give less weight to the between variation, compared to OLS.
- Two special cases can happen:
 - ① $\phi \rightarrow 0$: this means that σ_v is very small compared to σ_μ . In this case GLS converges to the within estimator.
 - ② $\phi \rightarrow 1$: this means that σ_v is very large compared to σ_μ . In this case GLS converges to the OLS (equal weight to within and between)

Fixed vs. random effects

- The individual effects are not fixed or random by nature and they can be modelled either as random or fixed depending on
 - ① the purpose of estimation
 - ② the probabilistic structure (assumptions)
 - ③ the correlation between the errors and the regressors
- in micro data, generally random effects are preferred (individuals are randomly drawn from a population), while in macro analysis, fixed effects are more popular (because the individual effects may be of interest per se)
- Assumption of uncorrelated effects, giving that $E(X^T v) = 0$.
Two situations:
 - ① $E(X^T \mu) = 0$: individual effects are not correlated with X . Both models (within and GLS) are consistent, but random effects (GLS) is more efficient than fixed effects (within)
 - ② $E(X^T \mu) \neq 0$: individual effects are correlated with X . Only the fixed effects (within) method gives consistent estimates as, with the within transformation, the individual effects vanish.

Application

Application in R: Example 2.2 (Croissant, Millo)

The two ways (error) model (i)

- The two ways error component is obtained by adding a time invariant effect λ_t to the model:

$$y_{nt} = \alpha + x_{nt}^T \beta + \mu_n + \lambda_t + v_{nt}$$

- Same assumptions hold for random time effects:
 - ① λ has zero mean and it is homoschedastic with variance σ_λ^2
 - ② time effects are mutually uncorrelated:
 $E(\lambda_t \lambda_s) = 0, \forall t \neq s$
 - ③ time effects are uncorrelated with the individual effects and with the idiosyncratic terms
- The error covariance matrix is
$$\Omega = \sigma_v^2 I_{NT} + \sigma_\mu^2 I_N \otimes J_T + \sigma_\lambda^2 J_N \otimes I_T$$

The two ways (error) model (ii)

The one-way
(error)
component
model

Ordinary least
squares

Generalized
least squares

Estimation
comparison

Two ways
error model

Advanced
arguments

References

- The error covariance matrix can be also expressed in terms of combination of idempotent and mutually orthogonal matrices:

$$\Omega = \sigma_v^2 W + (T\sigma_\mu^2 + \sigma_v^2)B_\mu + (N\sigma_\lambda^2 + \sigma_v^2)B_\lambda - \sigma_v^2 \bar{J}$$

where $B_\mu = I_N \otimes J_T / T$, $B_\lambda = J_T \otimes I_N / N$ and $\bar{J} = \frac{1}{NT} J_{NT}$

- $B_\mu \times x$ computes the individual means; $B_\lambda \times x$ computes the time means; $\bar{J} \times x$ computes the overall mean.
- The new within matrix W is $W = I_{NT} - B_\mu - B_\lambda + \bar{J}$

Fixed and random two ways models (i)

The one-way
(error)
component
model

Ordinary least
squares

Generalized
least squares

Estimation
comparison

Two ways
error model

Advanced
arguments

References

- As for the one way model, the two ways fixed effects model can be obtained in two different ways:
 - ① by estimating OLS on the model that includes individual and time dummies (two ways LSDV)
 - ② by estimating OLS on the model where all the variables have been transformed in deviation from time and individual means (pre multiplied by W , $z_{nt} - \bar{z}_n - \bar{z}_{.t} + \bar{\bar{z}}$)

Fixed and random two ways models (ii)

- For the GLS, similarly to what has been proposed in one way error models when NT is large, the variable are pre-multiplied by $\Omega^{-0.5}$, and then an OLS is applied on the transformed data
- Collecting terms, we obtain the following expression for the transformed data:

$$\tilde{z}_{nt} = z_{nt} - \theta_{\mu} \bar{z}_{n.} - \theta_{\lambda} \bar{z}_{.t} + (\theta_{\mu} + \theta_{\lambda} - \theta_2) \hat{\hat{z}}$$

where:

$$\begin{aligned} \textcircled{1} \quad \theta_{\mu} &= 1 - \frac{\sigma_v}{\sqrt{\sigma_v^2 + T\sigma_{\mu}^2}} \\ \textcircled{2} \quad \theta_{\lambda} &= 1 - \frac{\sigma_v}{\sqrt{\sigma_v^2 + N\sigma_{\lambda}^2}} \\ \textcircled{3} \quad \theta_2 &= 1 - \frac{\sigma_v}{\sqrt{\sigma_v^2 + T\sigma_{\mu}^2 + N\sigma_{\lambda}^2}} \end{aligned}$$

Application

Application in R: Example 2.8 and 2.9 (Croissant, Millo)

Advanced Arguments

- alternative ways to model the errors in error component models
- models to account for endogeneity
- models for unbalanced panels
- dynamic models accounting for time lagged variables
- model for count data
- models for spatial panels accounting for "space" lagged variables

Take home messages

- Different ways to account for heterogeneity (wipes out that or accounting for that)
- The individual and time effects may be modelled as fixed or random
- Within, Between, pooled OLS or GLS
- GLS and OLS gives intermediate results between within and between, with two extreme cases
- The use of fixed or random depends on the correlation assumption among the regressors and the errors

References

- Croissant, Y., & Millo, G. (2019). Panel data econometrics with R. Wiley
- Baltagi, B. (2008). Econometric analysis of panel data. John Wiley & Sons.